## Morus bassanus NORTHERN GANNET

Non-SPEC<sup>E</sup> (1994: 2) Status Secure Criteria — European IUCN Red List Category — Criteria — Global IUCN Red List Category — Criteria —

and the second	1
	and the second second
	in the second second
and the second	and the second s
and the second second	- FARA
	< 1
i and in the second secon	
and a state of the	A Providence of the
	Sector Sector Sector S
P Personal	C
- Busines	12 / 11 / 12 / 12 / 12 / 12 / 12 / 12 /
	all the state of the

Country	Breeding pop. size (pairs)	Year(s) Trend	Mag.%	References
Denmark				
Faroe Is.	2,000 - 2,000	95 (0)	(0-19)	
France	16,000 - 16,000	01 +	>80	6
Germany	70 - 70	95-99 +	50-79	
Iceland	23,000 - 25,000	84-94 +	0-19	7
Rep. Ireland	33,000 - 33,000	99-02 +	19	
Norway	4,200 - 4,200	02 +	20	35
Russia	50 - 60	90-00 +	N	80,154,155
UK	226,600 - 226,600	98-00 +	40	20
Total (approx.)	300,000 - 310,000	Overall trend Large increase		
Breeding range	>50,000 km <sup>2</sup>	Gen. length. 21	% Glo	bal pop. 75–94

Morus bassanus breeds at a relatively small number of sites in north-western Europe, which holds >75% of its global breeding population. Its European breeding population is large (>300,000 pairs), and increased substantially between 1970–1990. The species continued to increase across the majority of its European range during 1990-2000, and underwent a large increase overall. As a consequence of the range expansion accompanying its population growth, this previously Localised species is now evaluated as Secure.



2000 population	99		
1990 population	99		
	Data quality (%) – <i>Morus bassanus</i>		
1990-2000 trend	99		
1970–1990 trend	99		
2000 population	38 62		
1990 population	7 93		
	Data quality (%) – <i>Phalacrocorax carbo</i> (see p. 40, top) unknown poor medium good		
1990–2000 trend	37 61		
1970–1990 trend	20 80		